

## **Bipolarity, Choice, and Entropy**

Vladimir A. Lefebvre  
School of Social Sciences  
University of California at Irvine  
Irvine, CA 92697

[<valefebv@uci.edu>](mailto:valefebv@uci.edu)

February 20, 2004

## Abstract

Until now, formal models of bipolar choice have been phenomenological and were not related to the deep principles of processing information by live organisms, which limited the applications of these models and made it difficult to generalize them to the case of multi-alternative choice. We demonstrate here how to deduce a model theoretically based on a general definition of the self-reflexive system and one assumption which we called the Axiom of the Second Choice. We show further that such a deduction of the model reveals its unexpected connection to the relations between an internal variable of the self-reflexive system, a partial derivative of the entropy of the environmental influence, and a partial derivative of the entropy of choice made by the system. This connection allows us to expand the two-alternative model of bipolar choice to the case of an arbitrary number of alternatives.

## 1. Introduction

A real choice is often entailed with a conflict between profits and ideals. To analyze such situations, a special model of binary choice has been constructed; in it, the alternatives are polarized: one identifies the positive pole and the other identifies the negative one (Lefebvre, 1992; 2001). This model can be represented with the following equation:

$$\frac{1-X}{X} = \frac{1-x_1}{x_1} x_2, \quad (1)$$

where  $X$  is the probability of choosing the positive pole,  $x_1$ , where  $0 < x_1 < 1$ , is normalized attractiveness corresponding to the positive alternative, and  $x_2$ , where  $0 < x_2 < 1$ , is predetermined by factors lying beyond a given situation of choice. This parameter disturbs the proportionality between the probabilities of the alternatives' choice and their attractiveness, that is, the exact matching is broken. Although this model is widely used for simulation of decision-making processes (see for example, Adams-Webber, 1997; Kaiser & Schmidt, 2003; Miller, 2003; Miller & Sulcoski, 1999a,b; Taran, 1999; 2001), its applications are restricted by two factors. First, the model is phenomenological, that is, it is not based on the deep principles of the information processing by live organisms. Second, attempts to generalize this model to the multi-alternative choice were unsuccessful, because they were not justifiable. In this work we demonstrate that the model can be deduced from a very general definition of a self-reflexive system and an assumption which we called the Axiom of the Second Choice. It turns out that this model corresponds to the equation connecting an internal variable of the self-reflexive system with a partial derivative of the entropy of the environmental influence and with a partial derivative of the entropy of choice. This connection allows us to generalize the model to a case with  $n \geq 2$  alternatives.

## 2. Self-Reflexive System

A system is called *self-reflexive* if it can be presented with the following two functions:

$$B = \Phi_p(S) , \quad (2)$$

$$\Psi_p(B) = \Phi_p^{-1}(B) , \quad (3)$$

where  $B \in A_1$ ,  $p \in A_2$ , and  $S \in A_3$ . Variable  $B$  represents the behavior of the system, parameter  $p$  corresponds to the influence of the environment at the moment of choice, and  $S$  is an internal variable ('self') whose value does not depend on the environmental influence at the moment of choice.  $A_1$  is a set of the system's actions,  $A_2$  is a set of the environmental influences, and  $A_3$  is a set of values of the internal variable. Equation (2) corresponds to the statement that under fixed environment's influence, the behavior of the system may change only as a result of the internal variable change. Equation (3) shows that a reverse correlation must exist between the behavior and the internal variable (Lefebvre, 1965).

## 3. The Partial Derivative of Entropy

Consider Shannon's function

$$H = -\sum_{i=1}^n u_i \ln u_i , \quad 0 < u_i < 1, \quad u_1 + u_2 + \dots + u_n = 1 . \quad (4)$$

Variables  $u_1, u_2, \dots, u_i, \dots, u_{n-1}$  will be called *locally independent* if for any set of their values, a change of  $u_i$  for  $\Delta u_i$ , which is  $O(u_i)$ , does not change the value of  $u_j$  ( $j \neq i, j < n$ ), but changes only  $u_n$  for

$$\Delta u_n = -\Delta u_i .$$

In this case,

$$M_H(u_j) = \frac{\partial}{\partial u_j} H = -\sum_{i=1}^n \frac{\partial u_i}{\partial u_j} \ln(eu_i) = -\ln \frac{u_j}{u_n} , \quad j = 1, 2, \dots, n-1 , \quad (5)$$

where  $u_n = 1 - u_1 - u_2 - \dots - u_{n-1}$ .

## 4. A Model of Two-Alternative Bipolar Choice

While describing a model of the two-alternative bipolar choice we assume that the choice is made by a self-reflexive system. We assume also that the process of choice consists of two stages. First, the system makes a "preliminary" choice and then either realizes or cancels it. A choice which was realized will be called *actual choice*.

Let the system face a choice between positive and negative alternatives and let the probability that the positive alternative is chosen be given by

$$X(S) = \Phi_{x_1}(S) , \quad (6)$$

where  $0 < X < 1$ ,  $0 < x_1 < 1$ , and  $S \geq 0$ . Let

$$X(0) = x_1. \quad (7)$$

The value of  $x_1$  is the relative probability of the environmental push toward positive alternative. The numbers  $x_1$  and  $1-x_1$  are interpreted as normalized attractiveness of the positive and negative alternatives, respectively. Thus, equation (7) means that when  $S=0$ , the probability of the positive alternative being chosen is equal to its normalized attractiveness. Let us introduce the following assumption:

### ***Axiom of the Second Choice***

When the value of the internal variable grows from  $S$  to  $S + \Delta S$ , where  $\Delta S$  is  $o(S)$ , and the value of  $x_1$  remains unchangeable, the procedure of choice is as follows. First, the system performs a preliminary choice with the probability of the positive alternative being chosen is equal to  $X(S)$ . If it is chosen, the system realizes its choice. If the negative alternative is chosen, the system cancels its choice with probability  $c\Delta S$  (where  $c > 0$  and does not depend on  $S$ ) and repeats its choice again with the probability  $X(S)$  of choosing the positive alternative, after which realizes its choice independently of the chosen alternative's polarity.

It follows from this axiom that for the value of the internal variable equal to  $S+\Delta S$ , an actual choice is made with the probability of

$$X(S + \Delta S) = X(S) + (1 - X(S))X(S)c\Delta S , \quad (8)$$

from where

$$\Delta X(S) = c(1 - X(S))X(S)\Delta S .$$

When  $\Delta S \rightarrow 0$ , the equation for actual choice looks as follows:

$$\frac{dX(S)}{dS} = c(1 - X(S))X(S). \quad (9)$$

Equation (9) is well known and called *logistic* (see, for example, Mickens, 1990). By solving (9) under condition (7) we obtain

$$X(S) = \frac{x_1}{x_1 + (1 - x_1) \exp(-cS)} . \quad (10)$$

This expression can be represented as follows:

$$\frac{1-X}{X} = \frac{1-x_1}{x_1} \exp(-cS) . \quad (11)$$

We see that (1) is equivalent to (11), if  $x_2 = \exp(-cS)$ .

**Statement 1.** Reverse function of  $X = \Phi_{x_1}(S)$  exists and is given by

$$\Psi_{x_1}(X) = \frac{1}{c} (M_{H_1}(x_1) - M_{H_2}(X)) , \quad (12)$$

where

$$H_1 = -x_1 \ln x_1 - (1-x_1) \ln(1-x_1) ,$$

$$H_2 = -X \ln X - (1-X) \ln(1-X) ,$$

and  $0 < x_1 < 1$ ,  $0 < X < 1$ .

**Proof.** By substitution  $X(S)$  from (10) to (12) we obtain

$$\frac{1}{c} (M_{H_1}(x_1) - M_{H_2}(X(S))) = S . \quad (13)$$

We see that the self-reflexive system performing binary bipolar choice is described by equations (10) and (12) which correspond to general equations (2) and (3). Statement 1, in its essence, is a new theorem of the Information Theory. It establishes fundamental connections between the partial derivative of the entropy of the environment influence  $M_{H_1}(x_1)$ , the partial derivative of the entropy of choice  $M_{H_2}(X)$ , and the internal variable  $S$ .

## 5. A Model of Multi-Alternative Bipolar Choice

A connection of function  $\Psi_{x_1}(X)$  with the derivative of the entropy of choice allows us to extend the two-alternative model on the choice between any finite number of alternatives  $n \geq 2$ .

Let the self-reflexive system face a choice between  $n$  alternatives numbered  $i=1,2,\dots,n$ . Distribution of the probabilities of the environment influences is designated as  $q_1, q_2, \dots, q_n$  and called *normalized attractiveness*; distribution of the probabilities of choices is  $p_1, p_2, \dots, p_n$ . Let

$$H_1 = -\sum_{i=1}^n q_i \ln q_i \quad \text{and} \quad H_2 = -\sum_{i=1}^n p_i \ln p_i , \quad 0 < q_i < 1 \quad \text{and} \quad 0 < p_i < 1 .$$

Let alternatives 1, 2, ...,  $n-1$  be correlated with the positive pole, and alternative  $n$  with the

negative one. We will call alternatives  $1, 2, \dots, n-1$  *positive*, and alternative  $n$  *negative*. We assume that variables  $q_1, q_2, \dots, q_{n-1}$  as well as variables  $p_1, p_2, \dots, p_{n-1}$  are locally independent, that is, any small changes of their values can be compensated only by changes of variables  $q_n$  and  $p_n$  corresponding to the negative alternative. Therefore, in the generalized model, as well as in the two-alternative model, a change of a value belonging to the positive pole for  $\varepsilon$  leads to a change of a corresponding value belonging to the negative pole for  $-\varepsilon$ . The following system of equation is a generalization of (13):

$$\frac{1}{c}(M_{H_1}(q_j) - M_{H_2}(p_j)) = S_j, \quad S_j \geq 0, \quad j = 1, 2, \dots, n-1. \quad (14)$$

Consider each variable  $p_j$  where  $j=1,2,\dots, n-1$ , as an implicit function of independent variables  $q_1, q_2, \dots, q_{n-1}, S_1, S_2, \dots, S_{n-1}$ . By using (5) we can re-write (14) as

$$\frac{1}{c}(-\ln \frac{q_j}{q_n} + \ln \frac{p_j}{p_n}) = S_j, \quad j = 1, 2, \dots, n-1 \quad (15)$$

or

$$\frac{p_n}{p_j} = \frac{q_n}{q_j} \exp(-cS_j), \quad j = 1, 2, \dots, n-1, \quad (16)$$

where  $q_n = 1 - q_1 - q_2 - \dots - q_{n-1}$  and  $p_n = 1 - p_1 - p_2 - \dots - p_{n-1}$ .  
By solving system (16) we obtain

$$p_j = \frac{q_j}{\sum_{i=1}^{n-1} q_i \exp(c(S_i - S_j)) + q_n \exp(-cS_j)}, \quad j = 1, 2, \dots, n-1. \quad (17)$$

We will show now that the set of function (17) is compatible with the condition of local independence of variables  $p_j$ . Let us fix the values of all  $q_j$ . For any set of  $p_j$  values, due to (15), there is a set of  $S_j$  values which satisfies system (17). Therefore, for any  $O(p_n)$  change of  $p_j$  it is possible to find a set of  $S_1, S_2, \dots, S_{n-1}$  such that the values of  $p_i$  (where  $i \neq j$  and  $i < n$ ) do not change. Due to functions (15) continuity, small changes of  $p_i$  corresponds to small changes of  $S_j$ .

## 5. Integral Reflexive System

We will call a reflexive system *integral* in a case, in which all independent variables  $S_j$  take on the same values  $S \geq 0$ . When  $S=0$ , for all the alternatives the probabilities of choice are proportional to their normalized attractiveness; when  $S>0$ , the following equation holds:

$$\frac{p_i}{p_j} = \frac{q_i}{q_j}, \quad i, j = 1, 2, \dots, n-1, \quad (18)$$

i.e. only for positive alternatives are the probabilities of choice proportional to normalized attractiveness. For the ratio of the probability of choosing the negative alternative to the probability of choosing a positive one, this statement becomes wrong, which follows from (16).

The model can be extended to the arbitrary number of positive and negative alternatives under assumption that the system aggregates a set of negative alternatives into one negative alternative. After such aggregation, the model can predict the probability that an alternative from the set of negative alternatives will be chosen, but cannot predict the probability of choice for every negative alternative separately.

## Acknowledgment

I am grateful to my friends and colleagues, Jane Ryan, Stefan Schmidt, and Tim Kaiser for valuable advice. I also want to thank Victorina Lefebvre, without whose help this work would not be done.

## References

- Adams-Webber, J. (1997).  
Self-reflexion in Evaluating Others. *American Journal of Psychology*, **110**, 527-541.
- Kaiser, T. B. & Schmidt, S. E. (2003).  
Terrorist Recruitment and Reflexive Control. *Journal of Reflexive Processes and Control*, **2**, 2, 92-101.
- Lefebvre, V. A. (1965).  
On Self-reflexive and Self-organizing Systems, In: *Problemy Issledovania Sistem i Struktur*. Moscow: Izdatelstvo AN USSR. English translation in *Applied Ergonomics*, (1994), **1**, 6-9.
- Lefebvre, V. A. (1992).  
*A Psychological Theory of Bipolarity and Reflexivity*. Lewiston, N.Y.: The Edwin Mellen Press.
- Lefebvre, V. A. (2001).  
*Algebra of Conscience*, 2-nd enlarged edition. Dordrecht, Holland: Kluwer Academic Publishers.
- Lefebvre, V. A. (2003).  
Mentalism and Behaviorism: Merging? *Reflexive Processes and Control*, **2**, 2, 56-76.
- Mickens, R. E. (1990).  
*Difference Equation*. New York: Van Nostrand Reinhold.
- Miller, L. D. (2003).  
Essential Singularity Structure of Multi-Attribute Utility Function. *Reflexive Processes and Control*, **2**, 1, 103-118.
- Miller, L. D. & Sulcoski, M. F. (1999a).  
Reflexive Model of Human Behavior: Variations on Lefebvre's Theme. *Proceedings of the Workshop on Multi-Reflexive Models of Agent Behavior*. Los Alamos, NM: Army Research

- Laboratory, 51-62.
- Miller, L. D. & Sulcoski, M. F. (1999b).  
Application of Generalized Reflexive Behavior: Models for Situation-Specific Decisions. *Proceedings of the Workshop on Multi-Reflexive Models of Agent Behavior*. Los Alamos, NM: Army Research Laboratory, 69-94.
- Taran, T. A. (1999).  
The Formalization of Ethical Values in Models of Reflexive Behavior. *Proceedings of the Workshop on Multi-Reflexive Models of Agent Behavior*. Los Alamos, NM: Army Research Laboratory, 95-108.
- Taran, T. (2001).  
Many-Valued Boolean Model of the Reflexive Agent. *Journal of Multiple-Valued Logic*, 7, 97-127.