

Technical Report Series # MBS 09-07: **Long-Range Polymerase Chain Reaction Method for Detection of Human Red and Green Opsin Gene Polymorphisms**

Linda M. Wasserman, Monika K. Szeszel and Kimberly A. Jameson

Citation:

Wasserman, L.M., Szeszel, M. and Jameson, K.A. (2009). "Long-Range Polymerase Chain Reaction Analysis for Specifying Photopigment Opsin Gene Polymorphisms," Technical Report Series # MBS 09-07. Institute for Mathematical Behavioral Sciences University of California at Irvine, Irvine, CA, USA.

Author Notes:

A chronology of this research report is in order: The molecular genetic methods reported here were originally developed by L. M. Wasserman and K. A. Jameson from 1997-2000, and were subsequently refined between 2000 and 2002. The present methods extend the genetic analyses presented in Jameson et al. (2001) and permit greater specificity of the identified polymorphisms and more informative analyses of genotype-correlated behaviors.

Author contributions to this report are: (i.) method research and design (Wasserman and Jameson), (ii.) initial genotyping method implementation and assessment (Wasserman), (iii.) investigations confirming genotyping methods (Wasserman, Szeszel & Jameson), (iv.) data collection and statistical analyses of behavioral data (Jameson), and (v.) manuscript preparation (Jameson).

Writing of the present article was completed early in 2002 and until now was only available online in manuscript format. This IMBS Technical Report version of the paper is unchanged from the earlier 2002 version. In addition to the evaluation of the methods reported here, subsequent articles (referenced below) by Jameson, Sayim, Jameson, Szeszel and Alvarado (2005) and Jameson, Bimler and Wasserman (2006), further demonstrate the utility of these genotyping methods in analyzing color processing behaviors correlated with opsin genotypes. These genotyping procedures were also recently used to define some color vision observer variants that we mathematically model in evolutionary game theoretic simulations of color categorization (Jameson and Komarova 2009a, 2009b).

Sayim, B., Jameson, K.A., Alvarado, N. and Szeszel, M.K. (2005). "Semantic and Perceptual Representations of Color: Evidence of a Shared Color-Naming Function," *The Journal of Cognition & Culture* 5, 427-486.

Jameson, K.A., Bimler, D. and Wasserman, L.M. (2006). "Re-assessing Perceptual Diagnostics for Observers with Diverse Retinal Photopigment Genotypes," In *Progress in Colour Studies 2: Cognition*. Pitchford, N.J. and Biggam, C.P., eds. (Amsterdam: John Benjamins Publishing Co.), pp. 13-33.

Jameson, K.A. and Komarova, N. L. (2009a). "Evolutionary models of color categorization. I. Population categorization systems based on normal and dichromat Observers," *Journal of the Optical Society of America A*, 26(6), 1414-1423.

Jameson, K.A. and Komarova, N.L. (2009b). "Evolutionary models of color categorization. II. Realistic observer models and population heterogeneity," *Journal of the Optical Society of America A*, 26(6), 1424-1436.

Author Notes added by K. A. Jameson, October 2009.

Long-Range Polymerase Chain Reaction Method for Detection of Human Red and Green Opsin Gene Polymorphisms

Linda M. Wasserman

and

Monika K. Szeszel

Division of Medical Genetics

Department of Medicine and Cancer Center

University of California, San Diego

Kimberly A. Jameson*

Institute for Mathematical Behavioral
Sciences

University of California, Irvine

Corresponding author: IMBS, Social Science
Plaza, UC Irvine, Irvine CA, 92697-5100. Tel:
(949) 824-8651 Fax: (949) 824-3733
e-mail: kjameson@uci.edu

Human color-vision discriminates middle- from long-wavelength spectral light as a result of photopigment opsin genes found on the q-arm of the X-chromosome at location Xq28. These so called 'green' and 'red' opsin gene sequences are 98% homologous, producing middle-wavelength sensitive and long-wavelength sensitive photopigment opsin gene sequences which differ by only seven locations at which important amino-acid substitutions occur. Investigations of variations in green and red gene arrays, and the consequences on phenotype expression, are important for understanding expression mechanisms contributing to evolution of photopigment opsin genotypes. Here we introduce a long-range polymerase chain reaction assay for specifying the presence of genetic dimorphisms in human red and green gene sequences, and examine the utility of the long-range PCR in predicting perceptual behaviors found in phenotypes arising from identified genotypes. Consistent with behavioral differences correlated with results from a short-range PCR genotyping method (Jameson, Hightnote & Wasserman; 2001), the more specific long-range PCR analysis indicates that dimorphisms commonly found in green and red opsin genotypes are correlated with certain color perception behaviors. The long-range PCR presented provides an additional tool for use in evaluating hypotheses about photopigment opsin gene expression in color vision phenotypes.

Key words: X-linked photopigment opsin genes, Amino acid substitution, Spectral tuning, Color vision.

Introduction

Research on the genetic basis of retinal photopigments has enabled an understanding of the biochemical explanation for photopigment response sensitivity as well as the genetic basis for individual differences in color perception. Such work has shown that normal color vision in humans and old-world primates is trichromatic, being based on three classes of photopigments that are maximally sensitive to red (560–565

nm), green (530–535 nm), and blue (420– 430 nm) light (Dartnall et al. 1983; Schnapf et al. 1987, 1988; Bowmaker et al. 1978, 1991). The three photopigment opsins, as well as rhodopsin (the photopigment in retinal rods), are heptahelical proteins, composed of seven transmembrane α -helices that are linked by intra- and extracellular loops. Visual excitation following photon absorption occurs as the result of 11-*cis* to all-*trans* isomerization of the chromophore located at a binding site in helix 7.

The genes encoding the opsins or apoproteins of the human red and green photopigments are each composed of six exons and are arranged in a head-to-tail tandem array located on the q-arm of the X-chromosome (Nathans et al. 1986a,b; Vollrath et al. 1988; Feil et al., 1990). Individuals with normal color vision usually have one red opsin gene in the proximal position of the gene array and one or more green opsin genes. These X-linked opsin genes have 98% identity in nucleotide sequence (including introns and 3' flanking regions) (Zhao et al. 1990). The encoded "red" and "green" apoproteins differ by an estimated 15 residues, 7 of these known to occur at positions which influence photoreceptor responsivity in the expressed phenotype (Jacobs, 1998; Nathans et al. 1986a; reviewed in Nathans et al. 1992; Asenjo et al., 1994). The photopigment gene-specific amino acids are at codons 116, 180, 230, 233, 277, 285 and 309.

Human genotype/perceptual-phenotype analyses show that genotypic variation corresponds to shifts in the absorption spectra of expressed retinal pigments (Asenjo et al., 1994, Merbs & Nathans, 1992a, 1992b, 1993), with concomitant shifts in perceptual spectral sensitivity, or " λ -max," in human observers (Neitz, Neitz and Jacobs, 1991; Winderick et al. 1992, Neitz, Neitz and Jacobs, 1995).

More specifically, it has been shown that single amino-acid substitutions at codons 180 in exon 3 and codons 277 and 285 in exon 5 produce large shifts in phenotypic spectral sensitivity whereas the amino acids at codons 230 and 233 in exon 4 produce smaller shifts. (Merbs & Nathans, 1992a; Asenjo et al. 1994). The specific amino acids occurring at codons 180, 277 and 285 are highly conserved in vertebrates. The specific residues occurring at each position are associated with predictable shifts in λ -max for each species. Substitutions of amino acid residues involve the gain or loss of a hydroxyl-bearing group. Substitution of hydroxyl groups at key positions are associated with shifts in photopigment response sensitivity towards shorter wavelengths of the visible spectrum. Mammals whose MWS and LWS genes demonstrate codon conservation at these

key residues include cat, deer, guinea pig, horse, squirrel, goat, rabbit, dolphin, mouse, rat and several species of New World monkeys (Shyue et al, 1998; Yokoyama and Radlwimmer, 1999; Zhou et al, 1997).

Photopigments in invertebrates also consist of seven transmembrane α -helices. Although amino acid sequences in invertebrates are not conserved with respect to vertebrates, the specific amino acid residues at codons 180, 277 and 285 in red and green-sensitive photopigment genes are invariant in some insect species, e.g. *Papilio* (Briscoe, 2000), suggesting some evolutionary conservation of tertiary protein structure and biochemical mechanisms for photosignal transduction.

In contrast to the highly conserved relationship between the amino acids at codons 230, 233, 277 and 285 and perception of light as red or green, the specific amino acid occurring at codon 180 in each photopigment opsin gene is variable or polymorphic in *Homo sapiens* (Sharpe et al, 1998; Asenjo et al, 1994). In the Caucasian population, substitution of the amino acid serine for alanine in the MWS, or green, gene occurs in an estimated 6-9% of males. The amino acid alanine is substituted for serine in the LWS, or red, gene in an estimated 38% of individuals (Sharpe et al, 1998). Substitution of a hydrophobic residue for a hydroxyl-bearing amino acid at codon 180 produces a relatively large shift in λ -max in the LWS photopigment, but a lesser shift in λ -max in the MWS photopigment. Individuals with the more common serine at codon 180 in their LWS opsin gene will demonstrate an average spectral response, or λ -max, of 557 nm for red light. Individuals inheriting alanine at codon 180 in their LWS gene demonstrate a 5 nm shift in their average λ -max for red light to 552 nm, moving their spectral sensitivity for red light towards the λ -max for green light, which is 532 nm (Jacobs, 1998; Sharpe et al. 1998).

Complicating further the analysis of the relationship between genotype and perceptual behavior is the chromosomal location of the MWS and LWS genes. The MWS and LWS photopigment opsin genes occur in a tandem

array on the X chromosome. Females have two X chromosomes while males have a single X chromosome. Hence, females have two sets of such genes, one on each X chromosome, whereas males, who have only one X chromosome, have only one set of genes. As a result, females have potentially greater genetic variability in their MWS and LWS photopigment gene combination than is possible for males. Whereas there are four possible MWS/LWS genotype combinations at codon 180 for males, there are nine possible MWS/LWS genotype combinations for females. Thus, based on the reasoning that greater genotype diversity increases the diversity of expressed phenotypes, it might be reasonable to expect color perception variation in females not seen in male photopigment opsin genotypes.

In this article we present a new long-range PCR method (hereafter abbreviate LR PCR) for specifying amino-acid substitutions at exon 3, codon-180 of the MWS and LWS photopigment opsin genes. As described below, the method is a modification of existing methods using a short-range PCR technique which simply detects the presence of codon-180 polymorphisms (Jameson et al. 2001). The method described here extends the analyses of Jameson et al. by permitting greater specificity of the identified polymorphisms, and permits a more informative analysis of genotype-correlated behaviors reported by Jameson et al. (2001). In addition to predicting more accurately our measures of color perception behavior, the LR PCR is offered as a new tool to refine the correlation between perceptual behavior measures and photopigment opsin genotypes.

LR PCR Rationale and Method

Recent work examining the opsin genotype relationships to color vision behavior have frequently used genotyping analyses based on postmortem retinal mRNA (Hagstrom, Neitz and Neitz 1998; Balding, Sjoberg, Neitz and Neitz 1998). The present approach aims to supplement existing methods for the purpose of investigating the relationships among opsin genotype, expressed phenotype and color perception.

Because of the extensive DNA sequence homology between the green and red opsin genes, conventional PCR amplification of DNA through exon 3 has found it difficult to distinguish between MWS and LWS gene sequences at codon 180. The method to be described here uses a combination of molecular methods to enable accurate genotyping at exon 3, codon 180 for both the MWS and LWS genes. The method draws from the published work of Winderickx et al., Neitz and Neitz and colleagues, Asenjo et al. and Sharpe et al. In particular, the method is similar to one described by Winderickx et al. (1992) but differs because it incorporates an additional conformation of exon 4 and 5 gene-specific amino acids sequences. This additional confirmation is based on descriptions given by Asenjo et al. (1994) and Sharpe et al. (1998), which together distinguish the genomic regions of DNA sequence variation between MWS and LWS genes (Asenjo et al., 1994; Neitz, M. et al., 1995; Neitz, M. et al., 1991; Sharpe et al., 1998; Winderickx, et al. 1992). These studies provided the empirical justification for the method described here. The method uses a combination of three molecular approaches in order first to create MWS and LWS gene-specific DNA templates and then to use those templates to distinguish between their respective codon 180 sequences. A long-range polymerase chain reaction technique (LR PCR) generates gene-specific PCR products. DNA sequencing of each PCR template confirms this gene specificity, and PCR and a restriction digest determines MWS and LWS codon 180 genotypes.

The method enables accurate genotyping of the codon 180 polymorphism on each photopigment opsin gene. Thus fine comparisons can be made between genotype and color matching behavior. Analyses using this method demonstrate a close correlation with perceptual behavior and provide significant insight into mechanisms contributing to the variability in perceptual behavior.

Subjects. With permission of the UCSD Human Subjects Committee, informed consent was obtained from 38 female and 26 male UCSD undergraduates for participation in this

study. For simplicity of presentation, analyses in this article consider only data from female subjects. Three milliliters of venous blood from each student was collected into EDTA vacutainer tubes by a trained phlebotomist. Subjects were solicited through either the Psychology Department Human Subjects pool, or by posted solicitations for experimental participation for either cash payment for course extra-credit. To address specific empirical hypotheses subject solicitations were designed to maximize the yield of participants that were carriers or expressors of color vision deficiencies or anomalies. Thus, genotype frequencies of the present study do not represent population frequency estimates.

DNA Extraction Method. DNA was isolated from peripheral blood leukocytes using the PureGene DNA isolation kit (Gentra Systems, Minneapolis, MN).

Long Range PCR Rationale. As mentioned above, the genetic polymorphism of interest at codon 180 in the LWS and MWS genes occurs in exon 3 of each gene. Conventional PCR methods used to amplify DNA through this region of the gene cannot distinguish between DNA sequences within the LWS from DNA sequences within the MWS genes because the sequences of the two genes are identical except for the presence or absence of this polymorphism. The only significant region of sequence divergence between the two genes occurs in exon 5 which is more than 3000 bases away from the polymorphic region in exon 3 and beyond the limits of standard PCR methods. The LR PCR method described here makes use of PCR reagents specifically designed to extend the length of PCR-amplified DNA to several thousand base pairs. In addition, the method makes use of findings reported by Sharpe et al, (1998) of invariant amino acid sequences in exon 4 of each gene which distinguish LWS from MWS genomic DNA sequences. The method relies on these invariant sequences to confirm the gene specificity of each LR PCR product.

The LR PCR method consists of three steps:

(i.) Gene-specific antisense primers through exon 5 and a common sense primer recognizing sequences within intron 2 are used to amplify 4000bp (with 13646) base pair DNA products specific to each gene.

(ii.) A portion of each LR PCR product is amplified using primers through exon 4 to amplify and then DNA sequence PCR products which include codons 230 and 233 of each gene to confirm the gene specificity of each LR PCR product.

(iii.) Following confirmation of the gene-specificity of each LR PCR by DNA sequencing of exon 4, the remainder of each LR PCR product is amplified with primers from exon 3, followed by restriction digest, to determine the presence or absence of the codon 180 polymorphism.

Long Range PCR Technique. Following the manufacturer's recommended protocol, 100 ng DNA is added to a PCR mixture containing 1X Taq Extender buffer (Stratagene, La Jolla, CA), 350 uM each of dATP, dCTP, dGTP, and dTTP, 300 nM each primer, 5 U Taq Extender and 5 U Amplitaq DNA polymerase (Applied Biosystems, Foster City, CA) into a total volume of 28.6 ul. (Primer sequences are shown in Table 1 below.) Following denaturation at 94°C for 3 minute, each reaction undergoes 35 cycles of denaturation at 94°C for 30 seconds, annealing at 62°C for 30 seconds and extension at 72°C for 4 minutes, ending with a final 5 minutes of extension at 72°C. Below are the gene-specific sequences within exon 5 which were used to generate gene-specific exon 5 antisense primers. Unique amino acids are in italics and DNA bases unique to each gene are underlined and bolded.

Exon 4 Sequence Confirmation. 5 ul of LR PCR product is added to a PCR mixture containing 1.5 mM MgCl₂, 50 mM KCl, 10 mM TRIS HCl, pH 8.3, 0.001% gelatin, 200 uM each of dATP, dCTP, dGTP, and dTTP, 500 nM each

EXON 5

LWS Gene Codons:

274 275 276 277 278 279

Ile Phe Ala Tyr Cys Val
 DNA: **ATC TTT GCG TAC TGC GTC**

MWS Gene Codons:

274 275 276 277 278 279
Val Leu Ala Phe Cys Phe
 DNA: **GTC CTG GCA TTC TGC TTC**

primer (see Table 1 for primer sequences) and 1 U Taq DNA polymerase (Applied Biosystems) to a final volume of 50 ul. Following 3 minutes of denaturation at 94°C, each reaction undergoes 35 cycles of denaturation at 94°C for 1 minute, annealing at 60°C for 30 minute and extension at 72°C for 1 minute followed by a final extension at 72°C for 7 minutes. PCR products then undergo column purification (QIAGEN, La Valencia, CA) and are submitted to the UCSD Cancer Center DNA Sequencing Shared Resource for sequencing. Figure 1 below provides the gene-specific sequences within exon 4 used to confirm the specificity of each LR PCR product. Unique amino acids are in italics and DNA bases unique to each gene are bolded.

Figure 1. Gene-specific sequences within exon 4 of the red (LWS) and green (MWS) opsin genes. DNA sequence codings were used to confirm specificity of each long-range PCR product. Amino acids and DNA bases unique to each gene are shown in bold and italics.

EXON 4

230 231 232 233 234 235 236

LWS Gene
 Codons: *Ile* Pro Leu *Ala* Ile Ile *Met*
 DNA Sequence: **ATC** CCA CTC **GCT** ATC ATC **ATG**

MWS Gene
 Codons: Thr Pro Leu *Ser* Ile Ile *Val*
 DNA Sequence: ACC CCA CTC **AGC** ATC ATC **GTG**

Exon 3 Codon-180 Serine to Alanine

Polymorphism Detection. 5 ul LR PCR product is added to a PCR mixture containing 1.5 mM MgCl₂, 50 mM KCl, 10 mM TRIS HCl, pH 8.3, 0.001% gelatin, 200 uM each of dATP, dCTP, dGTP, and dTTP, 500 nM each primer and 1 U Taq DNA polymerase (Applied biosystems(as above)) to a final volume of 50 ul. Following 3 minutes of denaturation at 94°C, each reaction undergoes 35 cycles of denaturation at 94°C for 1 minute, annealing at 60°C for 30 seconds and extension at 72°C for 1 minute followed by a final extension at 72°C for 7 minutes. Digestion of 15 ul of exon 3 PCR product with 3 U of the restriction enzyme Fnu 4HI at 37°C for 1 hour determines the presence or absence of the codon 180 serine to alanine DNA polymorphism. When the digested PCR products are size-separated on a 3% agarose gel run in 1X TAE buffer, the presence of the DNA sequence coding for alanine is detected as a 152 bp band whereas the DNA sequence coding for serine is detected as a 193 bp band.




Table 1. Oligonucleotide primers used in amplification or sequencing reactions.

Long Range PCR Primers:	
Common Intron 2 Sense Primer	5'-GGC AAC ATA GTG AGA CCT CTT CTC -3'
LWS Exon 5 Antisense Primer	5'-CCAGCAGACGCAGTACGCAAAGAT-3'
MWS Exon5 Antisense Primer	5'-CCAGCAGAAGCAGAATGCCAGGAC-3'
Exon 4 Primers:	
Exon 4 Sense Primer	5'-ACAAACCCACCCGAGTTAG-3'
Exon 4 Antisense Primer	5'-GACTCATTTGAGGGCAGAGC-3'
Exon 3 Primers:	
Exon 3 Sense Primer	5'-TCATCTGTCTGCTCTCCCAT-3'
Exon 3 Antisense Primer	5'-ACCCTTACCTGCTCCAACCA-3'

Long Range PCR Results

Before describing results for this LR PCR genotyping method, a previously published genotyping method is summarized for comparison. Jameson, Highnote & Wasserman (2001) previously published a genotyping method for codon 180 dimorphisms which generates three possible genotypes in female subjects: Ser/Ser, Ser/Ala and Ala/Ala. Females who are Ser/Ser or Ala/Ala from this method have the same amino acid at codon 180 in both their LWS and MWS genes. For females who are Ser/Ala, it is not possible to determine which amino-acid is specific to LWS and MWS genotypes. Nevertheless, even with this restricted classification system, Jameson et al (2001) reasonably surmised that the "heterozygous" genotype individuals exhibiting *both* serine and alanine (i.e., Ser/Ala) necessarily possess a more *diverse* genotype than either homozygous genotype exhibiting a single amino-acid residue (i.e., Ser/Ser or Ala/Ala).

In light of the above mentioned fact that shifts in spectral sensitivity follow from codon-180 opsin gene substitutions, Jameson et al. (2001) tested the following hypothesis: *individuals with the genetic potential to express more diverse phenotypes (i.e., ser/ala) are likely to exhibit color perception behaviors which differ more than those in comparisons of homozygous genotypes (i.e., ser/ser or ala/ala)*. Using this simple genotyping method, Jameson et al. (2001) showed that individuals of the Ser/Ala classification demonstrated robust

differences in color perception behaviors when compared with their homozygous female counterparts. Results of the classification given by the Jameson et al. (2001) PCR method are summarized in Table 2, column 2.

Compared with the Jameson et al. method, under the new genotyping system which uses the LR PCR method described here, there are nine possible classifications of LWS-, MWS- genotypes at codon 180.

As illustrated by Figure 2's restriction gel digest and the LR PCR classification given in Table 2, column 1, accurate genotyping of codon 180 for both the LWS and MWS genes is achieved. Thus the LR PCR method successfully specifies the L- and M-cone genotypes of 37 female undergraduates. (Note hereafter N=37 due to insufficient DNA specimen for one participant).

Based on the results of the LR PCR, 37 female subjects can be classified into six of the nine possible genotype categories (Table 2, column 1). A comparison of columns 1 and 2 of Table 2 show that the LR PCR permits a more specific classification of subjects' opsin genotypes. The utility of the LR PCR's additional specificity can also be evaluated using the perceptual data of Jameson et al. (2001). For this we examine the degree of association between subject's perceptual data and the new LR PCR classification method, and compare that association with analogous measures under the original genotype classification method published by Jameson et al. (2001).




Figure 2. Fnu 4HI restriction digestion assay of genomic DNA from exon 3 codon-180 of the Green opsin (M-cone) gene of seven female donors (Lanes 1-7). Presence of the DNA sequence coding for alanine (i.e., denoted Ala.) is detected as a 160 bp band whereas the DNA sequence coding for serine (denoted Ser.) is detected as a 190 bp band. Lanes 1, 2 and 5, genomic DNA from human females with alanine at exon 3, codon-180 of the green gene. Lane 7, one female with serine at exon 3, codon-180. Lanes 3, 4, and 6, females with a serine-alanine dimorphism at exon 3, codon-180 of the green gene. Lane 8, DNA Ladder. Restriction gel digest products depicting analogous dimorphisms occurring at codon-180, exon 3 of the Red opsin (L-cone) gene are not depicted here, but are similar to those presented for the Green gene above.

Table 2. Frequencies of genotypes from a group of 37 participants evaluated using the new LR PCR method (column 1) compared with the original short-range PCR genotyping (column 2) from Jameson et al., (2001).

Long Range PCR Genotypes:	Original Genotyping from Jameson et al, 2001:
1. L-180-Ser/Ser & M-180-Ser/Ser : 10	Ser/Ser: 10
2. L-180-Ser/Ser & M-180-Ser/Ala : 7	Ser/Ala
3. L-180-Ser/Ala & M-180-Ser/Ala : 11	Ser/Ala
4. L-180-Ser/Ser & M-180-Ala/Ala : 1	Ser/Ala
5. L-180-Ser/Ala & M-180-Ala/Ala : 1	Ser/Ala
6. L-180-Ala/Ala & M-180-Ala/Ala : 7	Ala/Ala: 7

Relationships between LR PCR Genotype Classifications and Perceptual Behaviors

Through an additional step involving exon 4 sequence confirmation the LR PCR provides an accurate method for specifying L- and M-cone photopigment opsin genotypes. But the greater specificity illustrated in Table 2 is only of value in such studies if it reveals meaningful relationships in the genotype-phenotype linkage. Thus, for the Jameson et al. (2001) study, the question is whether the greater specificity provided by the LR PCR genotyping furthers our understanding of the genotype-phenotype linkage beyond the findings of the Jameson et al. (2001) classification system which showed that

a more diverse opsin genotype is associated with more complex (or richer) color experience behaviors.

Note that although the LR PCR genotyping provides greater specificity compared to the original genotyping method, it does not automatically follow that such specificity will necessarily be a better predictor of perceptual phenomena. With these issues in mind, we examine the perceptual results of the individuals classified by the two methods in Table 2 and ask the following two questions of the relationships in the data.

Table 3. Question 1 analyses using Kendall's *Tau* nonparametric measures of association between two genotyping classifications, O-PCR and LR PCR, and the Median Number of Chromatic Percepts, MNCP, experienced by 37 genotypically classified observers.

Variables correlated	Kendall's <i>Tau</i>	Significance level
O-PCR classification & MNCP	0.363	P < .002
LR PCR classification & MNCP	0.361	P < .0001

1. Which genotyping method is more informative regarding color-perception behaviors?

This question can be answered by measures of association between (i.) the two genotyping classifications [i.e., the *original PCR method* (abbreviated "O-PCR"), and the "LR PCR" method introduced above], and (ii.) the measures of perceptual behavior used in the Jameson et al. (2001) investigation. The measure of perceptual behavior is summarized here as "median number of perceived colors," and is tantamount to *the median number of different chromatic percepts a given observer detects in a series of judgments for diffracted spectrum stimulus* (see Jameson et al. 2001 for details of the stimulus and empirical task).

For the purpose of addressing the above question using correlational analyses, *Appendix A* provides the numeric code assigned to the O-PCR and LR PCR genotype classifications listed in Table 2. The coding system is an ordinal code following the hypothesis that greater genotype diversity predicts greater diversity in an observer's percept. For example, individuals genotyped as double-heterozygous for green and red gene dimorphisms (i.e., genotype in row 3, column 1, Table 2) are deemed the most genotypically diverse in our sample, and are assigned a ordinal code value of '4,' whereas genotypes with a lesser degree of genetic variation (i.e., homozygous genotype in row 1, column 1, Table 2) are assigned a lower ordinal value reflecting the likelihood of a *less-diverse* expressed phenotype. (See *Appendix A* for the O-PCR and LR PCR numeric codes.) Table 3 provides the Kendall's *Tau* statistics for the Question 1 analysis. Kendall's *Tau* is a nonparametric symmetric measure of ordinal association that ranges from '-1' (indicating a

total negative association between variables) to '1' (indicating total positive association).

Table 3's measures of association demonstrate that the different genotype classifications given by the original genotyping method (O-PCR) and the long-range genotyping method introduced here (LR PCR) are similarly correlated with the color perception behaviors observed using the Jameson et al. (2001) task. The measures indicate that in addition to giving a more specific genotyping, the LR PCR is at least as predictive of color perception behaviors as the originally used genotyping method.

One potential advantage of the LR PCR's genotyping specificity is that the additional information given by the classification permits evaluation of specific hypotheses related to the phenotypic expression of identified genes. For example, several competing hypotheses currently in the literature propose selective expression for M- versus L-genes which are distal from their locus of control region on the array (Yamaguchi, Motulsky & Deeb, 1997; Hayashi, Motulsky & Deeb, 1999; Shaaban & Deeb, 1998). One such hypothesis suggests that, despite the occasional presence of multiple red gene variants in the genetic array, expression mechanisms allow only a single red gene – that most proximal to the locus of control region – to be expressed phenotypically. Thus, it is believed that distal red gene variants do not phenotypically manifest (Nathans et al., 1989, 1993; Winderickx et al. 1992a). An alternative viewpoint suggests that multiple red genes can be phenotypically expressed (Sjoberg, Neitz, Balding & Neitz, 1998). Using the present LR PCR it is possible to examine plausible behavioral consequences of such hypotheses, as expressed in Question 2 below:

2. Under the assumption that multiple red gene variants are not expressed in the phenotype, which genotyping method is more informative regarding color-perception behaviors?

In essence, the assumption in Question 2 permits further assessment of Table 3's correlations between genotype classification and perceptual behavior. According to one theory at least, distal variants of red genes are not expressed. It follows from this idea that the perceptual consequences arising from the expression of green gene dimorphisms may be unchanged by the presence of more than one red gene variant (i.e., a L-opsin dimorphism) in the gene array.

As with Question 1, Question 2 is examined using Kendall's *Tau* measures. *Appendix A, section 2* describes a revised ordinal code which effectively lowers the rank assignment used to encode Table 2's, column 1 genotypes possessing more than one red gene variant. The basic relationship between the numeric values of the reassigned ordinal code and the genotype classifications remains preserved such that increasingly *diverse* photopigment opsin genotypes are encoded by monotonically increasing ordinal values.

Table 4's measures of association demonstrate that the genotype classification schemes given by the less-specific O-PCR method and the improved LR PCR method are also found to be well-correlated with differences in color perception behaviors similar to that seen in Table 3.

For Question 2's analysis the O-PCR numeric code remained unchanged from that described under the Question 1 analysis because the genotype classification of the O-PCR does

not permit further refinement. However, the LR PCR's specificity permits updating and, thus, Table 4's correlations are based on the recoded values of the LR PCR classifications that reflect increased association consistent with the hypothesis of increased diversity of the genotype, presumably because only one red gene is phenotypically expressed (i.e., Table 3, $\tau = 0.361$ versus Table 4, $\tau = 0.412$). This additional information permits analyses which reveal that the differences in color perception behaviors observed by Jameson et al. (2001) are likely attributable to increases in genotype dimorphisms. While this new information is consistent with some existing theories of opsin gene phenotype expression (Nathans et al., 1989, 1993; Winderickx et al. 1992a), direct tests are needed to determine whether the differences in perceptual behavior found by Jameson et al. (2001) for heterozygous females might be attributable to expressed variants of L-cone photoreceptor classes.

On this issue, however, it is informative that in analyses which the ordinal numeric code gives greater significance to green gene dimorphisms, the statistical trends show additional increases in the Kendall's *Tau* measures of association. These analyses suggest the LR PCR classification and the behavioral data are even more highly correlated. Although physiological confirmation is needed, such analyses underscore the suggestion that the LR PCR's specificity is informative concerning assessment of expressed phenotypes.

These correlational measures between the LR PCR and perceptual behaviors are a good indication that in addition to giving more specific genotyping classifications, the LR PCR is as useful a predictor of color perception behaviors as the short-range genotyping method used by Jameson et al. (2001). Moreover the LR

Table 4. Question 2 analyses using Kendall's *Tau* nonparametric measures of association between two genotyping classifications, O-PCR and LR PCR, and the Median Number of Chromatic Percepts, MNCP, experienced by 37 genotypically classified observers.

Variables correlated	Kendall's <i>Tau</i>	Significance level
O-PCR classification & MNCP	0.363	P < .002
LR PCR classification & MNCP	0.412	P < .0001

PCR's specificity permits the kind of hypothesis testing illustrated here in the straightforward assessment of the hypothesis of single red gene expression.

Summary

A long-range polymerase chain reaction technique for genotyping codon 180 polymorphisms in the MWS and LWS genes has been described. The method extends previous work using a short-range PCR technique (Jameson et al., 2001) and yields a classification of red and green photopigment opsin genotypes that permits detailed analysis of the perceptual consequences of the opsin gene polymorphisms. The utility of the technique was demonstrated through comparative analyses with a previously used genotyping method. In all analyses, the LR PCR method proved to be more accurate, more specific. The method served as a more sophisticated test of the hypothesis that more diverse opsin genotypes yield phenotypes producing more complex color perception experience. The LR PCR method also permitted evaluation of a hypothesis from the existing literature regarding the expression of gene variants located on the distal end of the genetic array. Although the ultimate proof of opsin gene expression mechanisms must be provided by physiologists and geneticists, this LR PCR method provides an alternative tool which might be used with existing methods for assessing the relationships between some opsin gene expression mechanisms, retinal photopigment phenotype, and color perception behaviors.

Author Notes

Supported by the National Science Foundation (NSF-9973903 to Jameson); a U.C.S.D. Hellman Faculty Fellowship Award (Jameson); and a U.C.S.D. Academic Senate Grant (#RZ649M to Wasserman). The authors gratefully acknowledge the U.C.S.D. Cancer Center's DNA sequencing shared resource.

References

- Asenjo, A.B. Rim, J. and Oprian, D.D. (1994). Molecular determinants of human red/green color discrimination. *Neuron*, 12, 1131-1138.
- Balding, S.D., Sjöberg, S.A., Neitz, J. and Neitz, M. (1998). Pigment gene expression in protan color vision defects. *Vision Research*, 38, 3359-3364.
- Bowmaker J.K., Dartnall H. J. A., Lythgoe J. N., Mollon J. D. (1978) The visual pigments of rods and cones in the rhesus monkey, *Macaca mulatta*. *Journal of Physiology: London*, 274, 329-348.
- Bowmaker JK, Astell S, Hunt DM, Mollon JD (1991) Photosensitive and photostable pigments in the retinae of Old World monkeys. *Journal of Experimental Biology*, 156, 1-19.
- Briscoe, A. D. (2000). Six opsins from the Butterfly *Papilio glaucus*: Molecular phylogenetic evidence for paralogous origins of red-sensitive visual pigments in insects. *Journal of Molecular Evolution*, 51, 110-121.
- Crandall, K. A., & Cronin, T. W. (1997). The molecular evolution of visual pigments of freshwater crayfishes (Decapoda: Cambaridae). *Journal of Molecular Evolution*, 45, 524-534.
- Dartnall HJA, Bowmaker JK, Mollon JD (1983) Human visual pigments: microspectrophotometric

results from the eyes of seven persons. *Proceedings of the Royal Society of London: Biology*, 220, 115–130.

Feil R., Aubourg P., Heilig R., Mandel J.L. (1990). A 195-kb cosmid walk encompassing the human Xq28 color vision pigment genes. *Genomics*, 6, 367–373.

Hagstrom, S. A., Neitz, J. and Neitz, M. (1998). Variations in cone populations for red-green color vision examined by analysis of mRNA. *NeuroReport*, 9, 1963-1967.

Hayashi, T., Motulsky, A.G. & Deeb, S.S. (1999). The molecular basis of deuteranomaly: position of a green-red hybrid gene in the visual pigment array determines colour vision phenotype. *Nature Genetics*, .

Jacobs, G.H. (1998). Photopigments and seeing: Lessons from natural experiments. *Investigative Ophthalmology and Visual Science*, 39, 2205-2216.

Jameson, K. A., Highnote, S. M. & Wasserman, L. M. (2001). Richer color experience in observers with multiple photopigment opsin genes. *Psychonomic Bulletin & Review*, 8, 244-261.

Meagher, M. J., Jorgensen, A. L., Deeb, S. S. (1996). Sequence and evolutionary history of the length polymorphism in intron 1 of the human red photopigment Gene. *Journal of Molecular Evolution*, 43, 622-630.

Merbs, S. L. & Nathans, J. (1992a). Absorption spectra of human cone pigments. *Nature*, 356, 433-435.

Merbs, S. L. & Nathans, J. (1992b). Absorption spectra of hybrid pigments responsible for anomalous color vision. *Science*, 258, 464-466.

Merbs, S. L. & Nathans, J. (1993). Role of hydroxyl-bearing amino acids in differentially tuning the absorption spectra of the human red and green cone pigments. *Photochemical Photobiology*, 58, 706- 710.

Nathans J, Thomas D, & Hogness DS. (1986a) Molecular genetics of human color vision: The genes encoding blue, green and red pigments. *Science*, 232, 193–202.

Nathans J, Thomas D, Hogness DS (1986b) Molecular genetics of inherited variation in human color vision. *Science*, 232, 203–210.

Nathans J, Merbs SL, Sung C-H, Weitz CL, & Wang, Y. (1992) Molecular genetics of human visual pigments. *Annual Review of Genetics*, 26, 401–422.

Neitz, M., Neitz, J. & Jacobs, G. H. (1995) Genetic basis of photopigment variations in human dichromats. *Vision Research*, 35, 2095-2103.

Neitz, M., Neitz, J. & Jacobs, G. H. (1991). Spectral tuning of pigments underlying red-green color vision. *Science*, 252, 971-974.

Neitz, M., Neitz, J. & Jacobs, G. H. (1995). Genetic basis of photopigment variations in human dichromats. *Vision Research*, 35, 2095-2130.

Schnapf J.L., Kraft T.W., & Baylor D.A. (1987) Spectral sensitivity of human cone photoreceptors. *Nature*, 325, 439–441.

Schnapf J.L., Kraft TW, Nunn BJ, & Baylor, DA (1988) Spectral sensitivity of primate photoreceptors. *Visual Neuroscience*, 1, 225–261.

Shaaban, S. A. & Deeb, S. S. (1998). Functional analysis of the promoters of the human red and green visual pigment genes. *Investigative Ophthalmology & Visual Science*, 39, 885-896.

Sharpe, L. T., Stockman, A., Jagle, H., Knau, H., Klausen, G., Reitner, A. and Nathans, J. (1998). Red, green, and red-green hybrid pigments in the human retina: Correlations between deduced protein sequences and psychophysically measured spectral sensitivities. *Journal of Neuroscience*, 18, 10053-10069.

Shyue, S.-K., Boissinot, S., Schneider, H., Sampaio, I., Schneider, M. P., Abee, C. R., Williams, L., Hewett-Emmett, D., Sperling, H. G., Cowing, J. A., Dulai, K. S., Hunt, D. M., & Li, W.-H. (1998). Molecular genetics of spectral tuning in new world monkey color vision. *Journal of Molecular Evolution*, 46, 697-702.

Sjoberg, S. A., Neitz, M, Balding, S. D., & Neitz, J. (1998). L-cone pigments genes expressed in normal colour vision. *Vision Research*, 18, 3213-3219.

Vollrath D, Nathans J, & Davies RW (1988)
Tandem array of human visual pigment genes at Xq28. *Science*, 240, 1669–1672.

Winderickx, J., Lindsey, D. T., Sanocki, E.,
Teller, D. Y., Motulsky, A. G., & Deeb, S. S. (1992).
Polymorphism in red photopigment underlies
variation in color matching. *Nature*, 356, 431-433.

Zhao, Z., Hewett-Emmett, D., & Li, W.-H.
(1998). Frequent gene conversion between human red
and green opsin genes. *Journal of Molecular
Evolution*, 46, 494-496.

Yamaguchi, T., Motulsky, A. G. & Deeb, S. S.
(1997). Visual pigment gene structure and expression
in human retinae. *Human Molecular Genetics*, 6,
981-990.

Yokoyama, S., & Radlwimmer, F. B. (1999).
The molecular genetics of red and green color vision
in mammals. *Genetics*, 153, 919-932.

Zhou, Y.-H., Hewett-Emmett, D., Ward, J. P. &
Li, W.-H. (1997). Unexpected conservation of the X-
linked color vision gene in nocturnal prosimians:
Evidence from two Bush Babies. *Journal of
Molecular Evolution*, 46, 494-496.

Appendix A: Ordinal codes assigned to genotype classifications used in Kendall's *Tau* correlational analyses of two empirical questions.

QUESTION 1: Which genotyping method is more informative regarding color-perception behaviors, the short-range PCR partitions from Jameson et al. (2001) or the new Long-Range PCR partitions?

Original-PCR Partition (or "O-PCR") Code from Jameson et al. (2001):

Below 'S' denotes serine, and 'A' denotes alanine, amino acids present at codon 180 in either, or both, Red and Green genes.

S/A: 2 (possibly heterozygous on one gene, possibly on both Red and Green genes)
 S: 1 (presumably homozygous-wildtype S & S on Red and green Genes)
 A: 1 (presumably homozygous-polymorphA & A on Red and Green genes)

Long-Range PCR (or "LR PCR") Partition Code:

S/A & S/A: 4 (double-heterozygous)
 S & S/A: 3 (homozygous-wild on Red, heterozygous on Green)
 S/A & A: 3 (heterozygous on Red, homozygous-wild on Green)
 S/A & S: 3 (heterozygous on Red, homozygous-polymorph on Green)
 A & S/A: 3 (homozygous-polymorph on Red, heterozygous on Green)
 S/S & A/A: 2 (multiple wildtype copies on Red, multiple wildtype copies on Green)
 S/S & S/S: 2 (multiple wildtype copies on Red, multiple polymorphic copies on Green)
 A/A & A/A: 2 (multiple polymorphic copies on Red, multiple wildtype copies on Green)
 S & A: 1 (homozygous-wild on Red, homozygous-wild on Green)
 A & S: 1 (homozygous-poly on Red, homozygous-poly on Green)
 S & S: 1 (homozygous-wild on Red, homozygous-poly on Green)
 A & A: 1 (homozygous-poly on Red, homozygous-wild on Green)

QUESTION 2: Under the assumption that multiple red gene variants are not expressed in the phenotype, which genotyping method is more informative regarding color-perception behaviors, the O-PCR partitions from Jameson et al. (2001) or the recoded LR PCR partitions?

Ordinal codes reassigned to LR PCR classification to reflect the theory that only one, proximal, red gene variant is expressed in the phenotype. In the New Long-Range PCR Partition code below, bolded code values reflect recoding based on this single red expression hypothesis. In the Question 2 analysis the O-PCR code remains as given above:

S/A & S/A: **3** (double-heterozygous)
 S & S/A: 3 (homozygous-wild on Red, heterozygous on Green)
 S/A & A: **2** (heterozygous on Red, homozygous-wild on Green)
 S/A & S: **2** (heterozygous on Red, homozygous-polymorph on Green)
 A & S/A: 3 (homozygous-polymorph on Red, heterozygous on Green)
 S/S & A/A: 2 (multiple wild copies on Red, multiple wild copies on Green)
 S/S & S/S: 2 (multiple wild copies on Red, multiple polymorphic copies on Green)
 A/A & A/A: 2 (multiple polymorphic copies on Red, multiple wild copies on Green)
 S & A: 1 (homozygous-wild on Red, homozygous-wild on Green)
 A & S: 1 (homozygous-poly on Red, homozygous-poly on Green)
 S & S: 1 (homozygous-wild on Red, homozygous-poly on Green)
 A & A: 1 (homozygous-poly on Red, homozygous-wild on Green).